

Stability of Total Phenolics and Anthocyanins of Three Types of Black Raspberry Confections During Storage


Aina Guo, Junnan Gu and Yael Vodovotz

Department of Food Science and Technology, The Ohio State University, 2015 Fyffe Ct., Columbus, OH 43210



Background

- ◆ Phenolic compounds, such as anthocyanins, proanthocyanidins and ellagitannins, are recognized as the major components in black raspberries that have been linked to the chronic disease prevention.
- ◆ Anthocyanin compounds are the common pigment in berries that give the berries blue to black colors based on the compositions and concentration^[1]. The anthocyanins in berries are related to the prevention of chronic disease, such as heart disease, cancer and obesity^[1].
- ◆ However, during long-term storage, the anthocyanins undergo the polymerization reactions^[1] to form colored polymeric compounds.^[2] The polymeric pigments have influence on the color and health-promotion ability related to anthocyanins and procyanidins.^[1]



Objective

The **objective** of this study was to determine the influence of storage temperature (4 °C and 25 °C) and storage time (two months) on the total phenolics, anthocyanins and percent polymeric color of three types of black raspberries confections: pectin gummy, starch gummy and hard candy.


Materials and Methods

Confection Preparation




- Each type of confections were prepared for three batches with the same freeze-dried black raspberry powder.
- Samples were stored at 4 °C and 25°C and phenolic compounds were measured during storage (fresh, 2, 4 6 and 8 weeks).

- ◆ Confections were cut and measured with different weight to make sure the amount of black raspberry powder is the same
- ◆ Measured confections were dissolved in 5% formic acid water
- ◆ Three different protocols were applied to the target compounds




Total Phenolics & Anthocyanins Measurements:

- Total Phenolics - **Folin-ciocalteu Colorimetry**^[3]





➤ Gallic acid (with concentration of 50, 100, 250 and 500 mg/L) served as calibration standards

- Anthocyanin - **pH-differential method**^[4]



- Percent polymeric color – **bisulfite**^[4]



Results

1. Total Phenolics during storage




Figure 1. Concentration of total phenolics in three types of confections during two-month storage.

2. Anthocyanins During Storage




Figure 2. Concentration of anthocyanins in three types of confections during two-month storage.

3. Polymeric Anthocyanins During Storage




Figure 3. Polymeric anthocyanins in three types of confections during two-month storage at 4 °C.




Figure 4. Polymeric anthocyanins in three types of confections during two-month storage at 25 °C.

Conclusions

- Total phenolics were relatively stable through the two-month storage at both 4 °C and 25 °C in pectin gummy and hard candy.
- Anthocyanins decreased during the first two-week storage and remained stable through the whole storage period at both temperature, except starch gummy stored at 25 °C, which have significant decrease after six week storage.
- Polymeric colors of all confections stored at room temperature increased during storage, while that confections stored at 4°C were more stable
- Phenolic compounds had the highest stability at 4 °C in pectin and hard candy confections during the two-month storage, while the starch gummy remained the highest concentration of total phenolics and anthocyanins

Reference

- [1] Howard L.R, Prior R.L, Liyanage R and Lay J.O. 2012. Processing and Storage Effect on Berry Polyphenols: Challenges and Implication for Bioactive Properties. J Agr Food Chem 60: 6678-93.
- [2] Kadivec M, Bornsek S.M, Polak T, Demsar L, Hribar J and Pozrl T. 2013. Phenolic Content of Strawberry Spreads during processing and Storage. J Agr Food Chem 61: 9220-9.
- [3] Waterhouse A.L. 2001. Determination of Total Phenolics. In Worstad R.E, editor. Current Protocols in Food Analytical Chemistry. New Jersey: John Wiley & Sons. I1.1.1-1.1.8
- [4] Giusti M.M and Wrolstad R.E. 2001. Characterization and Measurement of Anthocyanins by UV-Visible Spectroscopy. In Worstad R.E, editor. Current Protocols in Food Analytical Chemistry. New Jersey: John Wiley & Sons. F1.2.1-1.2.13

Acknowledgements

Thanks to Dr. Vodovotz, Junnan Gu and Jennifer Ahn-Jarvis for being great advisors and the Department of Food Science and Technology for opportunity to participate in research project.



THE OHIO STATE UNIVERSITY

COLLEGE OF FOOD, AGRICULTURAL,
AND ENVIRONMENTAL SCIENCES